



PRODUCT SPECIFICATION SHEET

Agar Medium C (Sabouraud-Dextrose Agar with Antibiotics) (DM1385I)

Intended Use

Agar Medium C (Sabouraud-Dextrose Agar with Antibiotics) (DM1385I) is recommended for selective cultivation and isolation of yeasts and moulds in compliance with IP.

Product Summary and Explanation

Sabouraud Dextrose Agar is a general-purpose medium devised by Sabouraud⁽¹⁾ for the cultivation of dermatophytes. The low pH of approximately 5.6 is favourable for the growth of fungi, especially dermatophytes, and slightly inhibitory to contaminating bacteria in clinical specimens.⁽²⁾ The addition of antibiotics is a modification designed to increase bacterial inhibition. Sabouraud Glucose Agar w/ Antibiotics is cited as Medium C and recommended for cultivation of yeasts and moulds by Indian Pharmacopoeia.⁽³⁾ This medium is a modification of Sabouraud Dextrose Agar originally formulated by Sabouraud.⁽¹⁾ The medium is used with antibiotics such as tetracycline and benzylpenicillin⁽⁴⁾ for the isolation of pathogenic fungi from materials containing large numbers of fungi or bacteria.

Principles of the Procedure

Agar Medium C (Sabouraud-Glucose Agar with Antibiotics) contains peptone which provides the nitrogen, amino acids and carbon. Dextrose monohydrate is an energy and carbon source. Tetracycline and benzyl penicillin inhibits a wide range of Gram-positive and Gram-negative bacteria, which makes the medium selective for fungi.⁽⁵⁾ The low pH favours fungal growth and inhibits contaminating bacteria from clinical specimen.

Formula / Liter

Ingredients	Gms / Liter
Dextrose monohydrate	40.00
Peptone (meat and casein)	10.00
Agar	15.00
Final pH : 5.6 ± 0.2 at 25°C	
Formula may be adjusted and/or supplemented as required to meet performance specifications	

Precautions

1. For Laboratory Use only.
2. IRRITANT. Irritating to eyes, respiratory system, and skin.
3. Some pathogenic fungi may produce infective spores, which are easily dispersed in air, so examination should be carried out in safety cabinet.

Directions

1. Suspend 61.36 grams of the medium in 995ml of purified/distilled water.
2. Heat to boiling to dissolve the medium completely.
3. Autoclave at 121°C, 15 psi pressure, for 15 minutes / validated cycle.
4. Aseptically add rehydrated contents of 1 vial of Tetracycline Selective Supplement (MS123).
5. Mix well and pour into sterile Petri plates.

Quality Control Specifications

Dehydrated Appearance	Cream to yellow homogeneous free flowing powder
Prepared Medium	Light amber coloured clear to slightly opalescent gel forms in Petri plates
Reaction of 6.14% solution	pH 5.6 ± 0.2 at 25°C
Gel Strength	Firm, comparable with 1.5% Agar gel





PRODUCT SPECIFICATION SHEET

Growth Promotion Test

Growth Promotion is carried out as per Indian Pharmacopoeia. Cultural characteristics observed with added Tetracycline Selective Supplement (FD196) after an incubation at 20-25°C for 48-72 hours (Incubate for 7 days for Trichophyton species)

Expected Cultural Response:

Sr. No.	Organisms	Results to be achieved				
		Inoculum (CFU)	Growth	Observed Lot value (CFU)	Recovery	Incubation Temperature/ period
1.	<i>Candida albicans</i> ATCC 10231	50 -100	good-luxuriant (white colonies)	25 -100	≥50 %	20 -25 °C/ ≤7 days
2.	<i>Aspergillus brasiliensis</i> ATCC 16404	50 -100	good-luxuriant	25 -100	≥50 %	20 -25 °C/ ≤7 days
3.	<i>Saccharomyces cerevisiae</i> ATCC 9763	50 -100	good-luxuriant	35 -100	≥50 %	20 -25 °C/ ≤7 days
4.	<i>Escherichia coli</i> ATCC 8739	≥10 ³	inhibited	0	0 %	20 -25 °C/ ≤7 days
5.	<i>Escherichia coli</i> NCTC 9002	≥10 ³	inhibited	0	0 %	20 -25 °C/ ≤7 days
6.	<i>Trichophyton rubrum</i> ATCC 28191	50 -100	good-luxuriant	25 -100	≥50 %	20 -25 °C/ ≤7 days
7.	<i>Lactobacillus casei</i> ATCC 334	≥10 ³	inhibited	0	0 %	20 -25 °C/ ≤7 days

The organisms listed are the minimum that should be used for quality control testing.

Test Procedure

Refer to appropriate references for standard test procedures.

Results

Refer to appropriate references and test procedures for interpretation of results.

Storage

Store the sealed bottle containing the dehydrated medium at 10 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedure

1. For identification, organisms must be in pure culture. Morphological, biochemical and/or serological tests should be performed for final identification.
2. Consult appropriate texts for detailed information and recommended procedures.





PRODUCT SPECIFICATION SHEET

Packaging

Product Name : Agar Medium C (Sabouraud-Glucose Agar with Antibiotics)

Product Code : DM1385I

Available Pack sizes : 500gm

References

1. Sabouraud K., 1892, Ann. Dermatol. Syphilol, 3:1061.
2. Murray, P. R 2005, In Manual of Clinical Microbiology, 7th ed., ASM, Washington, D.C.
3. Indian Pharmacopoeia 2010, Ministry of Health and Family welfare, Government of India, New Delhi
4. Ajello L., 1957, J. Chron. Dis., 5:545.
5. Lorian (Ed.),1980, Antibiotics In Laboratory Medicine, Williams and Wilkins, Baltimore.

Further Information

For further information please contact your local MICROMASTER Representative.





MICROMASTER LABORATORIES PRIVATE LIMITED

DM1385IPSS,QAD/FR/024,Rev.00/01.01.2018

Unit 38/39, Kalpataru Industrial Estate,
Off G.B. Road, Near 'R-Mall' , Thane (W) - 400607. M.S. INDIA.
Ph: +91-22-25895505, 4760, 4681. Cell: 9320126789.

Email: micromaster@micromasterlab.com

sales@micromasterlab.com

Prepared By	Checked By	Approved By
 01.01.2018	 01.01.2018	 01.01.2018
Microbiologist	Head Quality Control	Head Quality Assurance

Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.



Sabouraud Agar or Sabouraud Dextrose Agar (SDA) is a selective medium primarily used for the isolation of dermatophytes. Other fungi, yeasts, and filamentous bacteria such as *Nocardia* can also grow in SDA. The acidic pH of this medium (pH about 5.0) inhibits the growth of bacteria but permits the growth of yeasts and most filamentous fungi. Antibacterial agents can also be added to augment the antibacterial effect. Fungal colonies (front-side) on Sabouraud Dextrose Agar. This medium is also employed to determine the mycological evaluation of food, contamination in cosmetics, and clinically to aid in the diagnosis of yeast and fungal infections. High grade Agar Medium C (Sabouraud Glucose with Antibiotics) is supplied by Micromaster. It is available in the form of a cream to yellow homogenous free flowing powder with a pH in the range of 5.4 to 5.8. For maximum efficiency, it must be stored below 30°C in tightly closed container. Quantity provided: 500g /pack. Sabouraud Glucose with Antibiotics, also known as Medium C, is used for selective cultivation of yeasts and moulds, as per the British Pharmacopoeia. This medium is Carlier's modification of the formulation described by Sabouraud, for the cultivation of fungi. The medium is used with antibiotics such as Chloramphenicol for the isolation of pathogenic fungi from materials containing large numbers of fungi or bacteria. Sabouraud Dextrose Agar conforms with specifications of The United States Pharmacopeia (USP). Sabouraud Dextrose Agar is used in qualitative procedures for cultivation of pathogenic and nonpathogenic fungi, particularly dermatophytes. The medium is rendered more selective for fungi. 498. by the addition of antimicrobics. Sabouraud Dextrose Broth and Sabouraud Maltose Agar and Broth are also used for culturing yeasts, molds and aciduric microorganisms. Fluid Sabouraud Medium is used for cultivating yeasts, molds and aciduric microorganisms and for detecting yeasts and molds in normally sterile ... Sabouraud Glucose agar w/ antibiotics is cited as Medium C and recommended for cultivation of yeasts and moulds by British Pharmacopoeia (2). This medium was described originally by Sabouraud (7) for the cultivation of fungi particularly useful for the fungi associated with skin infections. The medium is used with antibiotics such as tetracycline and benzylpenicillin (1) for the isolation of pathogenic fungi from materials containing large numbers of fungi or bacteria. HMC peptone provides nitrogenous compounds. Glucose monohydrate provides an energy source.