

Stand structure, competition and growth of Scots pine (*Pinus sylvestris* L.) in a Mediterranean mountainous environment

Antonio GARCÍA-ABRIL, Susana MARTIN-FERNÁNDEZ, M. Angeles GRANDE, Jose A. MANZANERA*

Technical University of Madrid (UPM), Research Group for Sustainable Management, E.T.S.I. Montes, Ciudad Universitaria s.n., 28040 Madrid, Spain

(Received 19 January 2007; accepted 4 May 2007)

Abstract – The relationship between competition and tree growth was studied in four stands of *Pinus sylvestris* L. occurring in a continental Mediterranean mountain area (in the Guadarrama range, Spain), i.e., an uneven-aged stand, a stand with oak (*Quercus pyrenaica* Willd.) understorey, a plantation, and a mature even-aged stand. Competition was measured by a simple size-ratio distance-independent index and was negatively associated with tree diameter. This negative association was stronger in the uneven-aged, plantation and mature even-aged stands than in the stand with oak understorey. Competition was also negatively associated with current diameter increment. This relationship was moderately strong in the mature even-aged stand and weak in the uneven-aged stand and the plantation. In the uneven-aged and the mature even-aged stands, a weakly significant relationship was found between diameter growth and tree size, whereas these parameters were not associated in the stand with oak understorey. The competition index provided a better prediction of growth rate than the alternative use of diameter. Both diameter and basal area growth were greater in the uneven-aged than in the even-aged stands.

competition / growth / *Pinus sylvestris* / Scots pine / stand structure

Résumé – Structure des peuplements, compétition et croissance du pin sylvestre (*Pinus sylvestris* L.) dans un environnement montagneux méditerranéen. La relation entre compétition et croissance a été étudiée dans quatre peuplements de *Pinus sylvestris* L. que l'on rencontre dans la zone continentale des montagnes méditerranéennes (dans la région de Guadarrama en Espagne). Ont été pris en compte un peuplement inéquienne, un peuplement avec du chêne (*Quercus pyrenaica* Willd.) en sous-étage, une plantation, et un peuplement équienne âgé. La compétition a été mesurée par un index indépendant, simple rapport taille/distance, et était corrélée négativement avec le diamètre des arbres. Cette corrélation négative était plus forte dans le peuplement inéquienne, la plantation et le peuplement équienne âgé que dans le peuplement avec sous-étage de chêne. L'index de compétition était aussi corrélé négativement avec l'accroissement courant en diamètre. La relation était modérément forte dans le peuplement équienne âgé et faible dans le peuplement inéquienne et la plantation. Dans le peuplement inéquienne et le peuplement équienne âgé une relation faiblement significative a été trouvée entre croissance en diamètre et taille de l'arbre, alors que ces paramètres n'étaient pas corrélés dans le peuplement avec sous-étage de chêne. L'index de compétition fournit une meilleure prédiction du taux de croissance que le simple diamètre. La croissance en diamètre et en surface terrière était plus importante dans le peuplement inéquienne que dans le peuplement équienne âgé.

compétition / croissance / *pinus sylvestris* / pin sylvestre / structure de peuplement

1. INTRODUCTION

Scots pine (*Pinus sylvestris* L.) is one of the preferred species for reforestation in Spain, where it occupies third place among the native conifers (920 000 ha) and yields 7% of the total timber production. Average wood production is around 4 to 5 m³ ha⁻¹ year⁻¹, with wide site variations [4]. Such variability seems to be caused by structure, site quality, diseases and competition [7].

Scots pine is considered a shade-intolerant species, requiring much light for good development [23] and is unsuited to uneven-aged structure in Spain, although shading by retained trees can be helpful for regeneration [28], survival and growth [24]. In areas of continental climate, however, it behaves as a semi-shade-tolerant species and seedlings can tolerate some shading. Scots pine stands may be two-storied in sites with sufficient water supply. In some areas of Scotland, where there is no competition from other species and natu-

ral disasters such as forest fire are rare, regeneration in small gaps takes place [17]. As a result, trees of different age classes may be found simultaneously in the same stand [12]. This may lead to a clumped spatial structure, such as that found in some other pine species, for example *Pinus ponderosa* Douglas ex Lawson & C. Lawson [8], *Pinus nigra* Arn. [21] or *P. cembra* L. [24]. Currently, many forest managers actively pursue this structural variability because of the current social preference for diversity, whether in species, tree sizes or general stand structure [19].

There is, however, evidence that competition leads to reduced diameter increment in Scots pine stands [11]. Simple size ratio competition indices, such as that of Hegyi [9], explained over 50% of the variation in radial growth in Scots pine stands in Sweden [2].

The hypothesis of our work is that the current diameter growth of Scots pine is affected by competition in all types of stand structure. Therefore, the aim of this study is to characterise and compare competition-growth relationships in Scots pine stands of different structures.

* Corresponding author: joseantonio.manzanera@upm.es

Table I. Stand area, climatic, topographic and soil type data of the four Scots pine stands studied.

Stand	Stand area (ha)	P (mm)	PDP (mm)	DD (months)	Slope (%)	Soil type	Altitude (m)
Uneven-aged	98	1160.5	269.3	2.3	25	Haplustep	1450
With oak understorey	50	1328.5	319.3	1.9	16	Haplustep	1470
Plantation	10	1049.2	240.2	2.9	15	Haplustep	1420
Even-aged	12	1049.2	240.2	2.9	18	Haplustep	1495

P: annual precipitation; PDP: precipitation during the dry period (May–September); DD: drought duration.

2. MATERIAL AND METHODS

This study was conducted in three public forests in the Guadarrama mountains, located about 60 km from Madrid. There, four types of Scots pine stand structure were studied: a mature, naturally regenerated even-aged stand, an uneven-aged stand, a stand with oak understorey (*Quercus pyrenaica* Willd.) and a plantation. All four stands were situated at altitudes of between 1400 and 1500 m (Tab. I). Soils are formed on siliceous substrate of granites and gneiss [5], presenting umbric or ochric epipedon and a deep cambic horizon. They are classified as Haplusteps [27]. Mean annual temperature is 9.4 °C. Precipitation averages 1180 mm per year (Tab. I).

All the stands were first identified by photo-interpretation, visited by experts, and a representative area of each type of structure was delimited. In each of these representative areas of each stand structure two rectangular plots of 40 m by 60 m were chosen at random. In the plantation, only one plot was chosen given the smaller area and the greater homogeneity of the stand. Diameter at breast height (dbh, 1.3 m) and total height were measured in all trees higher than 1.3 m inside the plots. The homogeneity between both plots was verified and the data were merged as one sample per stand. A total of 1 209 trees were measured, 399 pine trees in the uneven-aged stand, 510 (251 oaks and 259 pine trees) in the stand with oak understorey, 170 pine trees in the plantation, and 130 pine trees in the even-aged stand.

For the variables age at breast height (1.3 m) and Current Diameter Increment in the last 5, 10 and 15 years (resp., CDI5, CDI10 and CDI15), a pilot sampling was previously carried out to keep sampling costs to a minimum. The measurement of these variables was restricted to Scots pine. Sample size for these variables was inferred assuming a sampling error of less than 9% as an objective, leading to a sampling size of 76 pine trees for the uneven-aged stand, 60 for the stand with oak understorey, 67 for the plantation and 41 for the even-aged stand. These pine trees were chosen by random sampling inside the plots. In each of these sampled trees, a wood core was extracted at a height of 1.3 m and annual radial increments were recorded. Also, the radius without bark was calculated for five-year periods. Current diameter increment (CDI) was estimated as twice the current radial increment; that is to say, twice the mean width of the last five (CDI5), 10 (CDI10) or 15 (CDI15) rings. Current basal area increment without bark was calculated as the difference between the present basal area and that of the previous five years, divided by five. The site index for Scots pine in the Guadarrama mountains (H_{100}) was determined with García and Gomez (1984) site index curves, based on dominant height (H_{dom}), average height of the 100 tallest trees per hectare, at 100 years age. Site index was estimated for the uneven-aged stands by deducing the mean age of the dominant trees from their wood cores and assuming that Scots pine frequently grows in clumps or copses of about 200 m² up to the age of 60 years or even more. These regeneration gaps behave like even-aged patches. The site index was then inferred from the García and Gomez site index curves (1984).

The effect of competition was described by Lorimer's index [14], as modified by Vayreda et al. [29]:

$$LM = \sum_j g_j/g_i$$

Where LM is the modified Lorimer area index; g_j is the basal area of competitor j ; $j = 1, \dots, n$; n is the number of competitors within a certain distance; g_i is the basal area of the subject tree. Distances between trees were calculated from relative Cartesian coordinates of each tree on the above-mentioned plots. A program was developed to calculate the basal area of the trees inside each neighbourhood radius and LM indices were calculated for increasing distances, m by m, from 1 to 10 m, and for 12.5 and 15 m.

For the variables dbh, CDI5, CDI10, CDI15 and LM indices at distances from 1 to 15 m, the relationship between pairs of variables was analysed, and the following regression models were fitted between each pair of variables: linear, double reciprocal, square root-Y, square root-X, exponential, S-curve, logarithmic, inverse-Y, inverse-X, multiplicative, logistic and log probit regression model. The model with the best fit was selected and the R -squared statistic was calculated for each model and stand.

3. RESULTS

Dominant height and stand density parameters are shown in Table II. Tables III to VI display the statistically significant R -squared (p -value < 0.01 for the hypothesis that R -squared = 0) between variables studied in the different stands. Age, height and diameter variables were strongly associated in both the uneven-aged stand and the stand with oak understorey, while these variables were only moderately associated in the plantation (data not shown). A range of ages was found in the plantation, as some retained trees were interspersed and natural regeneration from these older trees may have taken place. In the mature even-aged stand, dbh and height were strongly associated but the association between height and age was weaker, as all trees in this stand were from the dominant layer and variability was lower (data not shown).

Competition (LM index) was negatively associated with dbh in all stands, the effect being more pronounced at different distances among stands. In the uneven-aged stand, the R -squared was higher for 6 m distances or higher (more than 0.93 for dbh, Tab. III). In the stand with oak understorey, these relationships were stronger at neighbourhood radii of 10 m upwards, and ranging from a low R -squared of 0.06 for 1 m to an R -squared of 0.96 for 15 m (Tab. IV). In the plantation, dbh was strongly associated with competition at a distance of 6 m or more (R -squared > 0.8, Tab. V). In the mature even-aged stand (Tab. VI), dbh was strongly associated with competition,

Table II. Dominant height (H_{dom} : mean height of the 100 highest trees per ha), site index ($H_{100} = H_{dom}$ at 100 years age, [6]), mean age of the dominant trees, Stem number per ha (N), mean distance between trees ($MDBT = 100 N^{-1/2}$) in m, Relative Spacing ($RS = 100 (H_{dom} \times N^{1/2})^{-1}$) and basal area (g) in $m^2 ha^{-1}$ of four Scots pine stand structures located in the Guadarrama mountains.

Stand	H_{dom} (m)	H_{100} (m)	Mean dominant age (yr)	N (Stem No. ha^{-1})	$MDBT$ (m)	RS	g ($m^2 ha^{-1}$)
Uneven-aged	21	23	91	965	3.2	0.15	34.9
With oak understorey	19	17	124	1400*	2.7	0.15	32.9#
Plantation	22	29	64	703	3.8	0.17	59.0
Mature even-aged	24	25	98	315	5.6	0.23	55.4

* N includes 581 pines and 819 oaks ha^{-1} ; # basal area (g) of oaks was $0.93 m^2 ha^{-1}$.

Table III. Statistically significant relationships between diameter at breast height (dbh), current diameter increment of the last 5 (CDI5), 10 (CDI10) and 15 (CDI15) years and modified Lorimer's area indices for distances of 1 (LM1), 2 (LM2), 3 (LM3), 4 (LM4), 5 (LM5), 6 (LM6), 7 (LM7), 8 (LM8), 9 (LM9), 10 (LM10), 12.5 (LM12.5), and 15 (LM15) m in the uneven-aged stand.

	CDI5	CDI10	CDI15	LM1	LM2	LM3	LM4	LM5	LM6	LM7	LM8	LM9	LM10	LM12.5	LM15
dbh R -squared	0.13	0.17	0.18	0.17	0.42	0.58	0.53	0.68	0.94	0.96	0.97	0.98	0.99	0.99	0.99
p -value	0.0001	0.0002	0.0002	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001
CDI5 R -squared					0.13	0.12	0.13	0.16	0.31	0.29	0.34	0.31	0.28		
p -value					0.004	0.007	0.008	0.005	0.0001	0.0004	0.0002	0.002	0.008		
CDI10 R -squared					0.14	0.12	0.15	0.14	0.22	0.20	0.21				
p -value					0.0026	0.006	0.005	0.008	0.002	0.004	0.005				
CDI15 R -squared					0.15	0.15	0.19	0.21	0.24	0.24	0.25	0.23			
p -value					0.002	0.003	0.001	0.001	0.001	0.002	0.002	0.007			

Table IV. Statistically significant relationships between diameter at breast height (dbh), current diameter increment of the last five years (CDI5), and modified Lorimer's area indices for distances of 1 (LM1), 2 (LM2), 3 (LM3), 4 (LM4), 5 (LM5), 6 (LM6), 7 (LM7), 8 (LM8), 9 (LM9), 10 (LM10), 12.5 (LM12.5), and 15 (LM15) m in the stand with oak understorey.

	LM1	LM2	LM3	LM4	LM5	LM6	LM7	LM8	LM9	LM10	LM12.5	LM15
dbh R -squared	0.06	0.11	0.16	0.36	0.40	0.46	0.54	0.56	0.60	0.88	0.94	0.96
p -value	0.0005	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001
CDI5 R -squared											0.62	
p -value											0.0013	

Table V. Statistically significant relationships between diameter at breast height (dbh), current diameter increment of the last 5 (CDI5), 10 (CDI10) and 15 (CDI15) years and modified Lorimer's area indices for distances of 1 (LM1), 2 (LM2), 3 (LM3), 4 (LM4), 5 (LM5), 6 (LM6), 7 (LM7), 8 (LM8), 9 (LM9), 10 (LM10), 12.5 (LM12.5), and 15 (LM15) m in the plantation stand.

	CDI10	CDI15	LM1	LM2	LM3	LM4	LM5	LM6	LM7	LM8	LM9	LM10	LM12.5	LM15
dbh R -squared	0.10	0.19	0.12	0.43	0.50	0.47	0.75	0.83	0.89	0.93	0.94	0.93	0.97	0.99
p -value	0.008	0.0002	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001
CDI10 R -squared					0.14									
p -value					0.006									
CDI15 R -squared				0.15	0.24	0.18	0.20	0.26	0.20					
p -value				0.003	0.0003	0.003	0.002	0.0006	0.0067					

Table VI. Statistically significant relationships between diameter at breast height (dbh), current diameter increment of the last 5 (CDI5), 10 (CDI10) and 15 (CDI15) years and modified Lorimer's area indices for distances of 2 (LM2), 3 (LM3), 4 (LM4), 5 (LM5), 6 (LM6), 7 (LM7), 8 (LM8), 9 (LM9), 10 (LM10), 12.5 (LM12.5), and 15 (LM15) m in the mature even-aged stand.

	CDI5	CDI10	CDI15	LM2	LM3	LM4	LM5	LM6	LM7	LM8	LM9	LM10	LM12.5	LM15
dbh R -squared	0.22	0.25	0.34	0.34	0.28	0.38	0.67	0.82	0.89	0.88	0.91	0.90	0.95	0.99
p -value	0.0029	0.0011	0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001	< 0.0001
CDI5 R -squared						0.27	0.32	0.36	0.34	0.36	0.55	0.55		
p -value						0.0044	0.0029	0.002	0.005	0.0086	0.0011	0.0015		
CDI10 R -squared				0.36	0.35	0.31	0.45	0.55	0.51	0.54	0.61	0.59		
p -value				0.0004	0.0006	0.0022	0.0002	0.0001	0.0003	0.0006	0.0004	0.0008		
CDI15 R -squared				0.44	0.44	0.43	0.53	0.61	0.58	0.61	0.63	0.61		
p -value				0.0001	0.0001	0.0002	< 0.0001	< 0.0001	0.0001	0.0001	0.0003	0.0006		




Figure 1. Adjusted regression models between the current diameter increment of the last 15 years (mm year^{-1}) and the modified Lorimer's competition index at a distance of 7 m of the uneven-aged, mature even-aged and plantation stands. The real values in the scatter plot are represented by squares (\square) in the uneven-aged stand, stars (*) in the mature even-aged stand, and diamonds (\diamond) in the plantation.

with stronger coefficients for distances of 6 m or more (R -squared up to 0.99).

In the stand with oak understorey, no significant relationship was found between growth and size of pine trees (Tab. IV). In contrast, both in the uneven-aged and the mature even-aged stands, growth was significantly associated, although weakly, with size (Tabs. III and VI). In the plantation, only CDI_{10} and CDI_{15} were weakly associated with dbh (Tab. V).

Current diameter increment of the last 5, 10 and 15 years was negatively associated with the LM index for distances up to 10 m in the uneven-aged stand; a small amount of variation was explained by the model (R -squared up to 0.34 for CDI_5 at 8 m distance, Tab. III). No significant relationship was found between diameter growth and competition indices for longer distances. Similar results were obtained in the mature even-aged stand, with moderately stronger associations (R -squared up to 0.63 for CDI_{15} and LM_9 , Tab. VI). Competition indices up to a distance of 7 m were also significantly associated in the plantation with CDI_{15} (R -squared 0.26 for a 6 m distance, Tab. V), while CDI_{10} was only weakly associated with LM_3 (Tab. V). In contrast, growth showed no significant relationship with competition in the stand with oak understorey, except for CDI_5 and $LM_{12.5}$ (Tab. IV).


Regression models adjusted between current diameter increment (CDI) and competition index (LM) were compared for the uneven-aged, the even-aged and the plantation stands, i.e., CDI_{15} vs. LM_7 (Fig. 1). The stand with oak understorey had no significant model to be compared with other stands and was not included. In all cases, CDI of the uneven-aged stand is greater than that of the plantation and the even-aged stand. The same is true for the growth (i.e., CDI_{10}) vs. dbh model (Fig. 2). Furthermore, the estimated stand-level basal area in-

crement of the uneven-aged stand is greater than that of all the other stands (Tab. VII), although the basal area in the uneven-aged stand is lower than in the plantation and the mature even-aged stand (Tab. II).

4. DISCUSSION

Climate and site conditions of all four stands were similar (Tab. I), although site index (H_{100}) differences were observed between stands (Tab. II). This may influence growth, along with other forestry-related factors such as stand structure and competition. In other species, a relationship has been found between the radial increment of the stem and the site index [3, 18, 22]. Specifically, H_{100} of the stand with oak understorey was lower than H_{100} of the other stands, but dominant trees were also older than those in the other stands; this suggests a potential problem with the site index curves, the age measurement, or an interaction with the oak understorey.

As expected, a significantly negative relationship between competition and size was observed in all the stands. However, growth was in general weakly associated to size, with different patterns between the stands. The growth of larger and taller trees was faster in the uneven-aged stand, the plantation and the mature even-aged stand. This behaviour may be interpreted as a search for light and dominance by the larger trees following an asymmetric competition pattern [26]. In contrast, in the stand with oak understorey, growth was not associated with size. This may be attributed either to senescence of the larger pine trees or to symmetric competition, i.e., the effect of competition is the same on older, larger trees as on younger, smaller trees [26]. In this type of competition, the oak understorey may have played a role.



Stand	Regression model	R-squared	p
Uneven-aged	$CDI_{10} = \exp(1.26553 - 3.83567/dbh)$	0.17	0.0002
Mature even-aged	$CDI_{10} = 1/(0.305529 + 7.29771/dbh)$	0.25	0.0011
Plantation	$CDI_{10} = 1/(0.443136 + 12.7741/dbh)$	0.10	0.008

Figure 2. Adjusted regression models between the current diameter increment of the last 10 years (mm year^{-1}) and the diameter at breast height (dbh, cm) of the uneven-aged, mature even-aged and plantation stands. The real values in the scatter plot are represented by squares (\square) in the uneven-aged stand, stars (*) in the mature even-aged stand, and diamonds (\diamond) in the plantation.

Table VII. Current basal area increment without bark ($\text{m}^2 \text{ha}^{-1} \text{year}^{-1}$) in four stand structures. Current basal area increment was calculated as the difference between the present basal area and that of the previous five years, divided by five.

Stand structure	Current basal area increment ($\text{m}^2 \text{ha}^{-1} \text{year}^{-1}$)							TOTAL
	Diameter class (cm)							
	< 10 cm	10.1–20	20.1–30	30.1–40	40.1–50	50.1–60	> 60 cm	
Uneven-aged	0.09	0.11	0.08	0.12	0.19	0.08	0.01	0.68
With oak understorey	0.02	0.03	0.06	0.07	0.10	0.05		0.33
Plantation		0.01	0.17	0.26	0.13			0.57
Mature even-aged		0.00	0.02	0.05	0.08	0.13	0.07	0.35

Lorimer [14] reported that for predicting growth in natural even-aged hardwood stands distance-dependent measures of competition are not superior to distance-independent measures. Most of the comparisons between distance-dependent and distance-independent individual tree growth models do not report the expected differences in prediction ability [25]. Martin and Ek [16] also found the same result in uniform red pine plantations, concluding that distance-independent competition indices are more accurate for diameter growth projections in managed plantations, where location is of lesser importance. Vayreda et al. [29] compared several competition indices in two Scots pine stands in relation to the radial growth in the last five years and found that the modified Lorimer index performed better than all other indices tested. It is noteworthy that the Hegyi index, which takes into account distance to competitors, was no better than the modified Lorimer index, which does not include distance in its formula. Simple size ratio indices performed as well or better in their correlation with annual diameter growth than more complex indices, such as space, area overlap, and root/crown indices [10]. In con-

trast, Biging and Dobbertin [1, 2] obtained better results with estimated crown parameters. Our results show that the modified Lorimer index provides a substantially better prediction of growth rate than the alternative size parameters, such as dbh, in all the stand structures studied, especially in the mature even-aged stand.

The negative relationship between *CDI* and the *LM* index in all stands except for that with oak understorey may be interpreted as a hint of the negative effect of competition on growth when stand density is very high. At present, the low values of this relationship suggest that stand density is not so high as to cause serious growth limitation. The neighbourhood radius for maximum significant influence of competition on diameter growth was 8 m in the uneven-aged stand and 10 m in the mature even-aged stand. No relationship was found between *CDI5* and *LM* indices in the plantation, as a result of thinning during the last 10 years in this stand. In contrast, competition influenced Scots pine stem growth in other situations [15].

In our study, we observed a greater diameter growth in the uneven-aged stand than that in the mature even-aged stand and

the plantation for the same competition index. This result must be viewed with caution, given the small sample size. Nevertheless, other authors have reported that multiaged stands maintain comparable levels of timber productivity over time with even-aged stands [19], and Lähde et al. [13] observed higher productivity for uneven-sized stands of Scots pine in Finland. O'Hara and Nagel [20] also found that uneven-aged ponderosa pine stands were more efficient structures for maximizing the reception of sunlight than even-aged stands where the level of competition for available light and moisture was higher. In fact, even-aged stands had lower soil moisture availability per transpiration rate and a greater amount of water stress, which limits gas exchange and photosynthesis more than in multiaged structures [20].

5. CONCLUSIONS

The relationship between current diameter growth and competition was characterised in four stand structures of Scots pine, and an analysis was made of the variables influencing this relationship in a mountain area with a continental Mediterranean climate, i.e., in the Guadarrama range (Spain). The modified Lorimer competition index was successfully used to predict growth rate in both uneven-aged and even-aged stands. Taking into account the limited number of stands studied, no general conclusions can be drawn from the effect of stand structure on growth. Nevertheless, our results suggested a greater diameter and basal area growth of Scots pine in the uneven-aged stand than in the even-aged stand and the plantation. Additional research using this kind of approach might be warranted, and the effects on stand-level volume increment might also be considered. At densities that do not limit diameter growth, the uneven-aged stand maintains a continuous cover, a microclimate and is compatible with sustainable production. The conservation of a continuous cover and an irregular structure by means of moderate wood extractions for short periods is advisable for protection of the ecosystem and for aesthetic or social purposes.

Acknowledgements: This research work was supported by the D.G. de Investigación, Consejería de Educación y Cultura de la Comunidad de Madrid (project GR/AMB/0267/2004). We also thank Luis Orofino, Pablo Orofino, Jaime Hernandez, Alvaro Sanchez, Maria Estirado, Alvaro Ruiz, Elena Sanchez, Tania Lopez, Tania Caballero and Vanesa Jalvo for their assistance in field work and data processing, and to Pru Brooke-Turner for the linguistic revision of the manuscript.

REFERENCES

- [1] Biging G.S., Dobbertin M., A comparison of distance-dependent competition measures of height and basal area growth of individual conifer trees, *For. Sci.* 38 (1992) 695–720.
- [2] Biging G.S., Dobbertin M., Evaluation of competition indices in individual tree growth models, *For. Sci.* 41 (1995) 360–377.
- [3] Castedo-Dorado F., Dieguez-Aranda U., Alvarez-Gonzalez J.G., A growth model for *Pinus radiata* D. Don stands in northwestern Spain, *Ann. For. Sci.* 64 (2007) 609–619.
- [4] Diéguez-Aranda U., Álvarez-González J.G., Barrio-Anta M., Rojo-Alboreca A., Site quality equations for *Pinus sylvestris* L. plantations in Galicia (northwestern Spain), *Ann. For. Sci.* 62 (2005) 143–152.
- [5] Gandullo J.M., Nicolas A., Ecología de los pinares españoles. Ministerio de agricultura, I.F.I.E., Madrid, 1969.
- [6] García-Abejon J.L., Gomez-Loranca J.A., Tablas de producción de densidad variable para *Pinus sylvestris* L. en el Sistema Central, INIA, Madrid, 1984.
- [7] Gonzalez S.C., Bravo F., Inventario y descripción de la regeneración natural. Aplicación a grupos ordenados de pino silvestre (*Pinus sylvestris* L.) del Alto Ebro (Burgos), Montes 50 (1997) 21–28.
- [8] Harrod R.J., McRae B.H., Hartl W.E., Historical stand reconstruction in ponderosa pine forests to guide silvicultural prescriptions, *For. Ecol. Manage.* 114 (1999) 433–446.
- [9] Hegyi F., A simulation model for managing jack-pine stands, in: Fries J. (Ed.), Growth model for tree and stand simulation, Royal Coll. For., Stockholm, Sweden, 1974, pp. 74–90.
- [10] Holmes M.J., Reed D.D., Competition indices for mixed species Northern hardwoods, *For. Sci.* 37 (1991) 1338–1349.
- [11] Jähgagen K., Albrektsen A., Induced competition among Scots pine seedlings and its effect on future timber quality, *New For.* 12 (1996) 163–174.
- [12] Kuper J.H., Sustainable development of pine forests, Misc. Papers Wageningen University, 1994.
- [13] Lähde E., Laiho O., Norokorpi Y., Saksa T. Structure and yield of all-sized Scots pine dominated stands, *Ann. Sci. For.* 51 (1994) 111–120.
- [14] Lorimer C.G., Test of age-independent competition indices for individual trees in natural hardwood stands, *For. Ecol. Manage.* 6 (1983) 343–360.
- [15] Mäkinen H., Effect of intertree competition on biomass production of *Pinus sylvestris* (L.) half-sib families, *For. Ecol. Manage.* 86 (1996) 105–112.
- [16] Martin G.L., Ek A.R., A comparison of competition measures and growth models for predicting plantation red pine diameter and height growth, *For. Sci.* 300 (1984) 731–743.
- [17] Nixon C.J., Composition, structure and regeneration of tree species within Scotland's native pinewoods, in: Olsthoorn A.F.M. (Ed.), Management of mixed-species forests: silviculture and economics, IBN Scientific Contributions 15, IBN-DLO, Wageningen, 1999, pp. 191–198.
- [18] Nutto L., Spathelf P., Rogers R., Managing diameter growth and natural pruning of Parana pine, *Araucaria angustifolia* (Bert.) O Ktze., to produce high value timber, *Ann. For. Sci.* 62 (2005) 163–173.
- [19] O'Hara K.L., The Silviculture of transformation – a commentary, *For. Ecol. Manage.* 151 (2001) 81–86.
- [20] O'Hara K.L., Nagel L.M., A functional comparison of productivity in even-aged and multiaged stands: a synthesis for *Pinus ponderosa*, *For. Sci.* 52 (2006) 290–303.
- [21] Oldeman R.A.A., Forests: Elements of silvology, Springer, Heidelberg, 1990.
- [22] Sanchez-Gonzalez M., Tomé M., Montero G., Modelling height and diameter growth of dominant cork oak trees in Spain, *Ann. For. Sci.* 62 (2005) 633–643.
- [23] Schütz J.P., Sylviculture 1. Principes d'éducation des forêts, Presses Polytechniques et Universitaires Romandes, Lausanne, 1990.
- [24] Schütz J.P., Sylviculture 2. La gestion des forêts irrégulières et mélangées, Presses Polytechniques et Universitaires Romandes, Lausanne, 1997.
- [25] Soares P., Tomé M., Distance-dependent competition measures for eucalyptus plantations in Portugal, *Ann. For. Sci.* 56 (1999) 307–319.
- [26] Stoll P., Weiner J., Schmid B., Growth variation in a naturally established population of *Pinus sylvestris*, *Ecology* 75 (1994) 660–670.
- [27] USDA, Keys to soil taxonomy, 8th ed. 1998.
- [28] Valkonen S., Effect of retained Scots pine trees on regeneration, growth, form, and yield of forest stands, Investigación Agraria: Sistemas y Recursos Forestales: Fuera de serie 1 (2000) 121–145.
- [29] Vayreda J., Burriel J.A., Raventos J., Gracia C., Variabilidad en el crecimiento de *Pinus sylvestris* en función de su edad y tamaño y de la competencia local, *Studia Oecologica* 10/11 (1994) 373–384.

JAAghagen K., Albrektson A., Induced competition among Scots pine seedlings and its effect on future timber quality, *New For.* 12 (1996) 163-174. Kuper J.H., Sustainable development of pine forests, *Misc. Papers Wageningen University*, 1994. LAAhde E., Laiho O., Norokorpi Y., Saksa T. Structure and yield of all-sized Scots pine dominated stands, *Ann. Sci. For.* 51 (1994) 111-120 [CrossRef]. Stoll P., Weiner J., Schmid B., Growth variation in a naturally established population of *Pinus sylvestris*, *Ecology* 75 (1994) 660-670. USDA, Keys to soil taxonomy, 8th ed. 1998. Valkonen S., Effect of retained Scots pine trees on regeneration, growth, form, and yield of forest stands, *Investigaci3n Agraria: Sistemas y Recursos Forestales: Fuera de serie 1* (2000) 121-145. General Information. Common Name. Scots Pine. Scientific Name. *Pinus sylvestris*. Sun Tolerance. Full Sun. Height. Up to 35 m (up to 115 ft). Spread. 15 - 20 m (50 - 66 ft). Growth Rate. Fast. Bloom Time. Spring. Color. Green, Flower Color. Gold. Type. Tree. Native. USA, Europe. Classification. Kingdom. Plantae " Plants. Subkingdom. Tracheobionta " Vascular plants. Superdivision. Spermatophyta " Seed plants. Division. Coniferophyta " Conifers. Class. Pinopsida. Subclass. Order. Pinales. Family. Pinaceae " Pine family. Genus. Scotch pine (*Pinus sylvestris*), also called Scots pine, is an introduced species in North America, brought here from Europe probably in colonial days. Although it is used for both pulpwood and sawlogs, its principal value in the United States appears to be as a Christmas tree, as an ornamental, and for erosion control. The cool, humid climate of Scotland, along with the nature of the parent material, which is usually siliceous and acidic, frequently results in a deep litter and raw humus layer. The soils exhibit various degrees of podzolization. Scotch pine grows well on these soils but best growth is on freely drained sands and gravels, often on knolls and terraces. Aleppo pine forests consist of *Pinus halepensis* (Aleppo pine), which commonly colonises thermo- and calcicolous meso-Mediterranean scrub (EC 2007). In the Aegean islands, *Pinus pallasiana* occurs in mixed stands with *Pinus brutia* at higher areas of the broad *Pinus brutia* belt of Thasos, up to about 400 to 500 metres (Aegean pine forests of Thasos -subtype of the 9540 habitat) (EC 2007). 6. 91R0 Dinaric dolomite Scots pine forests (*Genisto januensis*-Pinetum). The 91R0 habitat consists of *Pinus sylvestris* woods of dolomitic soils of the Dinarides. These forests often occupy somewhat higher elevations than the similar dolomite *Pinus nigra* woods of the subtype Western Balkanic *Pinus nigra* forests (EC 2007). Ecological services and benefits o... Scots pine (*Pinus sylvestris* L.) is one of the preferred species for reforestation in Spain, where it occupies third place among the native conifers (920 000 ha) and yields 7% of the total timber production. Average wood production is around 4 to 5 m³ ha⁻¹ year⁻¹, with wide site variations [4]. Such variability seems to be caused by structure, site quality, diseases and competition [7]. Scots pine is considered a shade-intolerant species, requiring much light for good development [23] and is unsuited to uneven-aged structure in Spain, although shading by retained trees can be helpful for re...