



NORTHPARK Animal Hospital

FELINE ALLERGY

What are allergies and how do they affect cats?

One of the most common conditions affecting cats is allergy. An allergy occurs when the cat's immune system "overreacts" to foreign substances called allergens or antigens. Those overreactions are manifested in three ways. The most common is itching of the skin, either localized (one area) or generalized (all over the cat). Another manifestation involves the respiratory system and may result in coughing, sneezing, and/or wheezing. Sometimes, there may be an associated nasal or ocular (eye) discharge. The third manifestation involves the digestive system, resulting in vomiting or diarrhea.

Are there several types of allergies?

There are four known types of allergies in the cat: contact, flea, food, and inhalant. Each has common clinical signs and unique characteristics.

Contact Allergy

Contact allergies are the least common of the four types of allergies. They result in a local reaction to the skin. Examples of contact allergy include reactions to flea collars or to types of bedding, such as wool. If the cat is allergic to such substances, there will be skin irritation and itching at the points of contact. Removal of the contact irritant solves the problem. However, identifying the allergen can be challenging in many cases.

Flea Allergy

Flea allergy is the most common allergy in cats. A normal cat experiences only minor irritation in response to flea bites, often without any itching. The flea allergic cat, on the other hand, has a severe, itch-producing reaction when the flea's saliva is deposited in the skin. Just one bite causes such intense itching that the cat may severely scratch or chew itself, leading to the removal of large amounts of hair. There will often be open sores or scabs on the skin, causing a secondary bacterial skin infection. The area most commonly involved is over the rump or base of the tail. In addition, the cat may have numerous, small scabs around the head and neck. These scabs are called miliary lesions, a term which was coined because the scabs look like millet seeds.

The most important treatment for flea allergy is to get the cat away from all fleas. Therefore, strict flea control is the backbone of successful treatment. Unfortunately, this is not always possible in warm and humid climates, where a new population of fleas can hatch out every 14-21 days. However, the new topically applied monthly flea products may kill fleas before they have a chance to bite your cat. When strict flea control is not possible, injections of corticosteroids (or "cortisone" or "steroids") can be used to block the allergic reaction and give relief. This is often a necessary part of dealing with flea allergies. Fortunately, cats appear relatively more resistant to the side-effects of steroids than other species. If a secondary bacterial infection occurs, appropriate antibiotics must be used.

Inhalant Allergy

The most common type of allergy is the inhalant type, or atopy. Cats may be allergic to all of the same inhaled allergens that affect us. These include tree pollens (cedar, ash, oak, etc.), grass pollens (especially Bermuda), weed pollens (ragweed, etc.), molds, mildew, and the house dust mite. Many of these allergies occur seasonally, such as ragweed, cedar, and grass pollens. However, others are with us all the time, such as molds, mildew, and house dust mites. When humans inhale these allergens, we express the allergy as a respiratory problem; it is sometimes called "hayfever." The cat's reaction, however, usually produces severe, generalized itching. In fact, the most common cause of itching in the cat is inhalant allergy.

Most cats that have an inhalant allergy are allergic to several allergens. If the number is small and they are the seasonal types, itching may last for just a few weeks at a time during one or two periods of the year. If the number of allergens is large or they are present year-round, the cat may itch constantly.

Treatment depends largely on the length of the cat's allergy season. It involves two approaches. Steroids will dramatically block the allergic reaction in most cases. These may be given orally or by injection, depending on the circumstances. As stated previously, the side-effects of steroids are much less common in cats than in people. If steroids are appropriate for your cat, you will be instructed in their proper use.

Some cats are helped considerably by a hypoallergenic shampoo. It has been demonstrated that some allergens may be absorbed through the skin. Frequent bathing is thought to reduce the amount of antigen exposure through this route. In addition to removing surface antigen, bathing alone will provide some temporary relief from itching and may allow the use of a lower dose of steroids.

The second major form of allergy treatment is desensitization with specific antigen injections (or "allergy shots"). Once the specific sources of allergy are identified, very small amounts of the antigen are injected weekly. This is all in an attempt to reprogram the body's immune system. It is hoped that as time passes, the immune system will become less reactive to the problem-causing allergens. If desensitization appears to help the cat, injections will continue for several years. For most cats, a realistic goal is for the itching to be significantly reduced in severity; in some cats, itching may completely resolve. Steroids are not used with this treatment protocol, except on an intermittent basis. This therapeutic approach is recommended for the middle-aged or older cat that has year round itching caused by inhalant allergy. This approach is not used with food allergy.

Although desensitization is the ideal way to treat inhalant allergy, it does have some drawbacks and may not be the best choice in certain circumstances.

1. **Cost:** This is the most expensive form of treatment.
2. **Age of Patient:** Because many cats develop additional allergies as they get older, young cats may need to be retested 1-3 years later.
3. **Success Rate:** About 50% of cats will have an excellent response. About 25% get partial to good response. About 25% get little or no response. The same statistics are true for people undergoing desensitization.
4. **Food Allergies:** Although tests for food allergy are available, the reliability of the test is so low that it is not recommended at this time. A food trial remains the best diagnostic test for food allergy.
5. **Time of Response:** The time until apparent response may be 2-5 months, or longer.
6. **Interference of steroids:** Cats must not receive oral steroids for two weeks or injectable steroids for six weeks prior to testing; these drugs will interfere with the test results.

Food Allergy

Cats are not likely to be born with food allergies. More commonly, they develop allergies to food products they have eaten for a long time. The allergy most frequently develops in response to the protein component of the food; for example, beef, pork, chicken, or turkey. Food allergy may produce any of the clinical signs previously discussed, including itching, digestive disorders, and respiratory distress. We recommend testing for food allergy when the clinical signs have been present for several months, when the cat has a poor response to steroids, or when a very young cat itches without other apparent causes of allergy. Testing is done with a special hypoallergenic diet. Because it takes at least eight weeks for all other food products to get out of the system, the cat must eat the special diet exclusively for 8-12 weeks (or more). If positive response occurs, you will be instructed on how to proceed. *If the diet is not fed exclusively, it will not be a meaningful test.* We cannot overemphasize this. If any type of table food, treats or vitamins are given, these must be discontinued during the testing period.

Because cats that are being tested for inhalant allergy generally itch year round, a food allergy dietary test can be performed while the inhalant test and antigen preparation are occurring.

INSTRUCTIONS

Those instructions, which are specific for your cat, have been checked

- ___ 1) An injection of steroids was given. Relief should be apparent within 12-24 hours. If not, please call. The cat should feel better and itch less for about one month. If an increase in water consumption or urination occurs, please report this to us for future reference. These side-effects are common with steroid administration and will go away in a few days without treatment. Return for further evaluation when the first signs of itching recur.
- ___ 2) Begin oral steroids when the first signs of itching return. Give ___ tablets every other evening. Adjust dosage upward or downward to the lowest effective dose, with a maximum dose of _____ permitted. Stop giving the medication every 4-6 months to see if there are times of the year when therapy is not needed. Report any increase in water consumption to us at once.
- ___ 3) We have dispensed oral steroid tablets. Prednisone is most commonly used. The specific drug being dispensed for your cat is labeled on the bottle. Give ___ of the ___ mg tablets every other day for 3 doses (6 days), then ___ tablets every other day for 3 more doses (6 more days). Continue this downward progression (_____) until the first signs of itching recur. At that time, go back to the next higher level and report that level to us (so we can be sure it is a safe level). Stop giving the tablets every 4-6 months to see if there are periods of the year when they are not necessary. When itching returns, begin immediately at the maintenance dose. If that does not stop the itching, increase the dosage slightly (to a maximum of ___ tablets) for a few doses, then return to the lower dose. Report any increase in water consumption to us at once. (This may occur at the initial dose but should stop on the maintenance dose.)
- ___ 4) Your cat is to exclusively eat a hypoallergenic diet. If it will not do so readily, mix it 25:75 with the current diet for several days, then gradually increase the special diet to 100%. If this does not work, contact us for an alternative plan. Discontinue any chewable treats or vitamins. Table food is not allowed. If possible, offer only distilled water to drink.
- ___ 5) Your cat has a flea allergy or has enough fleas to make the other allergy problem worse. Flea control is very important and should include treating the cat and its environment. Bear in mind that flea allergies often accompany other types of allergies, especially inhalant allergy.
- ___ 6) Your cat has a bacterial skin infection secondary to allergy. The following are recommended:
- a) Antibiotics are to be used for the next _____ days. If the infection is not gone by the time the medication is completed, call for a refill or for a change in medication.
 - b) The medicated shampoo, _____, is to be used every _____ days. Allow the shampoo to stay in the haircoat for a few minutes before thoroughly rinsing the cat.
 - c) The topical medication, _____, is to be used _____ times daily for _____ days.

Cats exposed to allergens may develop allergies or allergic reactions. Allergies tend to become evident and intensify over extended periods of time and can take years to develop. Some allergic diseases and allergies in cats include feline atopic dermatitis, flea allergy dermatitis, feline-mosquito hypersensitivity, and food-induced allergy. In the case of feline atopy, hypersensitivity to allergens is due to genetic predisposition. However, various allergies may arise due to environmental factors... "Feline food allergies often cause skin problems. Gastrointestinal symptoms are less common. Before treatment other common allergies need to be eliminated from consideration such as those to fleas and mites. Treatment involves eliminating most food groups down to a single protein to see if your cat gets better. Food groups will be gradually reintroduced until the problem food is identified." Cat food allergies usually cause skin reactions. In some cases symptoms include diarrhea and vomiting. Feline Allergies in Cats. a.k. Atopy, inhaled allergies, contact allergies. Show table of contents. Written by. It's important to note that in most cases the diagnosis of allergic disease is arrived at via exclusion, meaning that other causes of similar symptoms are ruled out, making the diagnosis of allergic disease the most likely.