

## Log $P$ —Making Sense of the Value

Sanjivanjit K. Bhal

Advanced Chemistry Development, Inc.

Toronto, ON, Canada

[www.acdlabs.com](http://www.acdlabs.com)

### Introduction

The fact that water and many organic substances do not mix but form separate layers when combined together has far-reaching implications for chemistry, biology, and the environment. The partition coefficient ( $P$ ) describes the propensity of a neutral (uncharged) compound to dissolve in an immiscible biphasic system of lipid (fats, oils, organic solvents) and water. In simple terms, it measures how much of a solute dissolves in the water portion versus an organic portion. Solutes that are predominantly dissolved in the water layer are called hydrophilic (water liking) and those predominantly dissolved in lipids are lipophilic (lipid liking). The partition coefficient is an important measurement of the physical nature of a substance and thereby a predictor of its behavior in different environments. The log $P$  value provides indications on whether a substance will be absorbed by plants, animals, humans, or other living tissue; or be easily carried away and disseminated by water.<sup>1</sup>

As a result of its wide and varied applications, the partition coefficient is also referred to as Kow or Pow.

The log $P$  value is a constant defined in the following manner:

$$\text{Log}P = \log_{10} (\text{Partition Coefficient}) \quad \text{Partition Coefficient, } P = [\text{organic}]/[\text{aqueous}]$$

Where [ ] indicates the concentration of solute in the organic and aqueous partition.

A negative value for log $P$  means the compound has a higher affinity for the aqueous phase (it is more hydrophilic); when log $P$  = 0 the compound is equally partitioned between the lipid and aqueous phases; a positive value for log $P$  denotes a higher concentration in the lipid phase (i.e., the compound is more lipophilic). Log $P$  = 1 means there is a 10:1 partitioning in Organic : Aqueous phases.

Although log $P$  is a constant, its value is dependent on the choice of the organic partitioning solvent and, to a lesser degree, on the conditions of measurement. ACD/Labs' log $P$  algorithms specifically calculate partitioning between octan-1-ol and water—the most commonly used system.

In this application note, we will discuss specific applications of log $P$  in order to offer a deeper understanding of this parameter and to demonstrate its wide-ranging impact in chemistry.

### Applications of Log $P$

As a fundamental property of matter, lipophilicity is a descriptor that can help scientists predict and understand the transport and impact of chemicals in physiological and ecological systems. Log $P$  values are important to many industries and areas of research in determining how to deliver chemical substances to specific sites, or eliminate chemicals from others, as well as limiting unwanted dispersal of chemicals through the environment.

## Drug Discovery

$\log P$  is used in the pharmaceutical/biotech industries to understand the behavior of drug molecules in the body. Drug candidates are often screened according to  $\log P$ , among other criteria, to help guide drug selection and analog optimization. This is because lipophilicity is a major determining factor in a compound's absorption, distribution in the body, penetration across vital membranes and biological barriers, metabolism and excretion (ADME properties). According to 'Lipinski's Rule of 5' (developed at Pfizer) the  $\log P$  of a compound intended for oral administration should be  $<5$ . A more lipophilic compound:

- Will have low aqueous solubility, compromising bioavailability. If an adequate concentration of a drug cannot be reached or maintained, even the most potent in-vitro substance cannot be an effective drug.
- May be sequestered by fatty tissue and therefore difficult to excrete; in turn leading to accumulation that will impact the systemic toxicity of the substance.
- May not be ideal for penetration through certain barriers. A drug targeting the central nervous system (CNS) should ideally have a  $\log P$  value around 2;<sup>2</sup> for oral and intestinal absorption the ideal value is 1.35–1.8, while a drug intended for sub-lingual absorption should have a  $\log P$  value  $>5$ .

Not only does  $\log P$  help predict the likely transport of a compound around the body. It also affects formulation, dosing, drug clearance, and toxicity. Though it is not the only determining factor in these issues, it plays a critical role in helping scientists limit the liabilities of new drug candidates.

## Agrochemicals

Use of  $\log P$  in insecticide, fungicide, and herbicide research is in many ways similar to that in drug discovery. It helps scientists to understand the transport and loss process of chemicals. In agrochemical discovery and development, scientists are charged with finding chemicals that are effective for a specific action while being relatively harmless to surrounding plants, insects, humans, and the environment (especially waterways).

Bioavailability in agrochemistry refers to uptake of a chemical by a plant through soil or foliage, or ingestion by a pest. 'Briggs Rule of 3' suggests that for an agrochemical to be bioavailable (among other parameters) it should have a  $\log P$  value  $<3$ .

Other reasons to achieve an appropriate balance between hydrophilicity and lipophilicity by understanding and modifying  $\log P$  are:

- While the best mode of application of a chemical may be spraying as an aqueous solution, the active compound must be lipophilic enough to resist leaching into waterways. For chemicals that end up in the water table, environmental impact must also be considered.
- Toxicity of agrochemicals must be understood and is also related to  $\log P$ . Wildlife such as fish and birds are obviously open to more direct exposure, but since chemicals can carry over into food by being taken up by the plant, human toxicity must also be understood.

## Environmental Chemistry

The environmental impact of chemicals has become a more visible issue over the years. Concerns include pharmaceutical and agrochemical residues, and waste chemicals generated by industry.  $\log P$  is one molecular property factor that can help in developing an environmental profile of the fate of chemicals from bioaccumulation in soil, exposure to wildlife, and the impact to human health.

## Flavors and Fragrances

New flavors and fragrances are a powerful driving force behind food, beverages, cosmetics, and household products. We come into contact with (or ingest) many products from these categories on a daily basis and lipophilicity has some interesting implications.

Partitioning between lipid and aqueous media is said to influence flavor perception.<sup>3</sup> Achieving an acceptable  $\log P$  value for the fragrance in fabric softener is important for deposition of perfume on the fabric after rinsing. As well as helping to predict the desired effect of a compound, understanding lipophilicity helps scientists determine their in-vivo fate (ADME properties and toxicity) in case of ingestion.

This handful of uses are not the only areas for which  $\log P$  is important. For example, there is on-going research on  $\log P$  as one important parameter in helping to classify the irritant and/or corrosive nature of chemicals in order to minimize the number of animals required in the testing of new substances.<sup>4</sup>

## How Can I Obtain $\log P$ Values?

The partition coefficient of a compound can be experimentally measured using a variety of widely accepted methods. The two most common methods are Shake Flask and HPLC. The Shake Flask Method, suitable for the broadest range of solutes, is rather time-consuming, though generally thought to be the most accurate. High performance liquid chromatography (HPLC) is a faster method that can be used when the chemical structure of the solute is known.

Prediction of  $\log P$  is another method for obtaining this information. Software, such as ACD/LogP,<sup>5</sup> uses algorithms to calculate the  $\log P$  of a compound by the sum of its fragments ( $\log P$  values for individual fragments are obtained from experimental data and/or determined statistically). Prediction is a powerful resource because it can be done in the absence of a compound sample. It is used, even by groups that carry out measurements of  $\log P$ , to plan experiments and verify results.

## Conclusion

$\log P$  is an important molecular physical property that impacts a wide range of systems. Used in conjunction with other critical parameters, it can help drive research forward in many industries and help determine the ultimate fate of chemical substances in our environment. Although there are well-established methods for measurement of the partition coefficient, these can be cumbersome and costly. Using prediction of  $\log P$  before a substance is even synthesized offers a means to guide scientists toward more fruitful research and development.

## References

1. A. Leo, C. Hansch, D. Elkins. (1971). Partition Coefficients and their uses. *Chemical Reviews*, 6: 525–616.
2. C. Hansch, J. P. Björkrot, A. Leo. (1987). Hydrophobicity and Central Nervous System Agents: On the Principle of Minimal Hydrophobicity in Drug Design. *J. Pharm. Sci.*, 76(9): 663–687.
3. M. J. Greenberg. (1980). The importance of hydrophobic properties of organic compounds on their taste intensities: a quantitative structure-taste-intensity study. *J. Agric. Food Chem.*, 28(3): 562–566.
4. P. Ruiz, G. Begliutti, T. Tincher, J. Wheeler, M. Mumtaz. (2012). Prediction of Acute Mammalian Toxicity Using QSAR Methods: A Case Study of Sulfur Mustard and Its Breakdown Products. *Molecules*, 17(8): 8982–9001.
5. ACD/LogP, [www.acdlabs.com/logp/](http://www.acdlabs.com/logp/) Advanced Chemistry Development, Inc., Toronto, ON, Canada.

LogP—Making Sense of the Value. Sanjivanjit K. Bhal. Advanced Chemistry Development, Inc. Although logP is a constant, its value is dependent on the choice of the organic partitioning solvent and, to a lesser degree, on the conditions of measurement. ACD/Labs™ logP algorithms specifically calculate partitioning between octan-1-ol and water—the most commonly used system. In this application note, we will discuss specific applications of logP in order to offer a deeper understanding of this parameter and to demonstrate its wide-ranging impact in chemistry. LogP values are important to many industries and areas of research in determining how to deliver chemical substances to specific sites, or eliminate chemicals from others, as well as limiting unwanted dispersal of chemicals through the environment. Some of these applications are discussed below. Application Note Drug Discovery LogP is used in the pharmaceutical/biotech industries to understand the behavior of drug molecules in the body. Drug candidates are often screened according to logP, among other criteria, to help guide drug selection and analog optimization. This is because lipophilicity is a major determining factor in a compound's absorption, distribution in the body, penetration across vital membranes and biological barriers, metabolism and excretion (ADME properties). Sensemaking or sense-making is the process by which people give meaning to their collective experiences. It has been defined as "the ongoing retrospective development of plausible images that rationalize what people are doing" (Weick, Sutcliffe, & Obstfeld, 2005, p. 409). The concept was introduced to organizational studies by Karl E. Weick in the 1970s and has affected both theory and practice. Weick intended to encourage a shift away from the traditional focus of organization theorists on decision... AppNote LogP - Making Sense of the Value. Learn how logP is measured, what the values mean, and how logP is applied in various R&D industries. Read more. AppNote Lipophilicity Descriptors: Understanding When to Use LogP & LogD. It represents the tendency of a compound to differentially dissolve in these two immiscible phases (typically, Octanol and Water). The partition coefficient is also referred to as Kow and the octanol/water partition coefficient. LogP prediction models estimate this value as a logarithmic ratio (logP, ClogP, or AlogP). The partition coefficient acts as a quantitative descriptor of the lipophilicity (or hydrophobicity) of a compound. What is the difference between logP and logD? Blog titles: Making sense of value: what is value? The value index: how to measure value? How to balance benefits delivery with risk optimisation to realise value. These have also been published as a collective piece. Case studies. The Value Management Toolkit and Transport in Leeds, David Worsley LinkedIn. Using Value Engineering to guide organisational design studies, Behnam Bashokoo LinkedIn. Companies, small and large, are continuously faced with the need to realign themselves to ever-changing markets, competitions and ways of working.