

## ANALYSIS OF BIOMARKERS FROM EDIACARAN FOSSILS: BRINGING TOGETHER PALAEOLOGY AND ORGANIC GEOCHEMISTRY

Ilya Bobrovskiy, Janet M. Hope, Jochen J. Brocks

Australian National University, Australia

The emergence of modern animal phyla in the Cambrian was precluded by the appearance of large architecturally complex organisms, the Ediacara biota (571 to 541 Ma). These fossils might hold clues to the understanding of history of the animal lineage and the evolution of complex life on Earth. However, most of them have evaded taxonomic classification, with interpretations ranging from marine animals or giant single-celled protists to bacterial colonies and terrestrial lichens (Xiao and Laflamme, 2009).

A technique of analysing biomarkers from individual fossils opens a new dimension in the study of these enigmatic fossils. Instead of looking at their morphology, it allows to analyse remains of organic molecules produced by these organisms. Biomarkers represent skeletons of biomolecules produced by all living organisms; as different groups of organisms possess different lipids, biomarkers can be used to unravel their biological origins (Brocks and Pearson, 2005; Volkman, 2005). Thus, based on morphology, Ediacaran fossils *Beltanelliformis* have been variably interpreted as bacterial colonies, planktonic algae, benthic algae, fungal colonies, jellyfish bodies, sponge-like animals and sedentary coral polyps. Biomarker analysis showed that *Beltanelliformis* contain abundant hopanes and long-chain *n*-alkanes with odd-over-even predominance, indicating that these fossils represent large spherical colonies of cyanobacteria similar to modern *Nostoc* (Bobrovskiy et al., 2018b). Biomarkers from the most iconic Ediacaran macrofossil *Dickinsonia*, along with two other dickinsoniid genera, *Andiva* and *Yorgia*, indicate that they were not lichens or giant protists, but in fact belong among the oldest animals preserved in the rock record (Bobrovskiy et al., 2018a).

The proposed technique has broader applications than uncovering the biological origins of ancient organisms. In addition to sterols they produce, animals may contain molecules from what they have eaten. Indeed, using biomarkers, we were able to detect the presence of a gut in some of Ediacaran macroorganisms and distinguish the lipid composition of their last meal. Gut content analysis is one of the most powerful tools for deciphering trophic structure of ancient ecosystems and ecology of their members (Butterfield, 2001; Vannier, 2012). Biomarkers provide a new dimension to these studies, allowing to analyse the gut content of ancient macroorganisms even when it is not fossilized.

### References

- Bobrovskiy, I., Hope, J.M., Ivantsov, A., Nettersheim, B.J., Hallmann, C., Brocks, J.J., 2018a. Ancient steroids establish the Ediacaran fossil *Dickinsonia* as one of the earliest animals. *Science* 361, 1246-1249.
- Bobrovskiy, I., Hope, J.M., Krasnova, A., Ivantsov, A., Brocks, J.J., 2018b. Molecular fossils from organically preserved Ediacara biota reveal cyanobacterial origin for *Beltanelliformis*. *Nature Ecology & Evolution*.

- Brocks, J.J., Pearson, A., 2005. Building the Biomarker Tree of Life. *Reviews in Mineralogy and Geochemistry* 59, 233-258.
- Butterfield, N.J., 2001. Cambrian Food Webs, in: Briggs, D.E., Crowther, P.R. (Eds.), *Palaeobiology II*. Blackwell Scientific, Oxford, pp. 40-43.
- Vannier, J., 2012. Gut Contents as Direct Indicators for Trophic Relationships in the Cambrian Marine Ecosystem. *PLOS ONE* 7, e52200.
- Volkman, J.K., 2005. Sterols and other triterpenoids: source specificity and evolution of biosynthetic pathways. *Organic Geochemistry* 36, 139-159.
- Xiao, S., Laflamme, M., 2009. On the eve of animal radiation: phylogeny, ecology and evolution of the Ediacara biota. *Trends Ecol Evol* 24, 31-40.

Analysis of Biomarkers from Ediacaran Fossils: Bringing Together Palaeontology and Organic Geochemistry. By Ilya Bobrovskiy, Janet M. Hope and Jochen J. Brocks. *Analysis of Biomarkers from Ediacaran Fossils: Bringing Together Palaeontology and Organic Geochemistry*, Page 1 of 1. < Previous page | Next page > /docserver/preview/fulltext/2214-4609/2019/P166-1.gif. There is no abstract available for this article. Molecular fossils from organically preserved Ediacara biota reveal cyanobacterial origin for Beltanelliformis. *Nature Ecology & Evolution*. [Google Scholar]. Brocks, J., Pearson, D., 2005. Building the Biomarker Tree of Life. *Reviews in Mineralogy and Geochemistry* 59, 233–258. [Google Scholar]. Butterfield Charnia at 50: developmental analysis of Ediacaran fronds. *Palaeontology*, 51, 11–26. Antcliffe, J. B. and McCloughlin, N. 2008. Deciphering fossil evidence for the origin of life and the origin of animals: common challenges in different worlds. 211–229. In Seckbach, J. and Walsh, M. (eds). *Dickinsonia from Ediacara: a new look at morphology and body construction*. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 270, 311–323. Brasier, M. D. and Antcliffe, J. B. 2009. *Organic Geochemistry*, 30, 1411–1428. Hoffman, P. F. and Schrag, D. P. 2002. The snowball Earth hypothesis: testing the limits of global change. Biomarkers, molecular clocks, and microRNAs suggest a 200-Myr missing Precambrian fossil record of siliceous sponge spicules. *Geobiology*, 8, 24–36. The Ediacaran (Ediacaran; formerly Vendian) biota is a taxonomic period classification that consists of all life forms that were present on Earth during the Ediacaran Period (c. 635–541 Mya). These were composed of enigmatic tubular and frond-shaped, mostly sessile, organisms. Trace fossils of these organisms have been found worldwide, and represent the earliest known complex multicellular organisms. PDF | On Sep 4, 2019, Ilya Bobrovskiy and others published Analysis of Biomarkers from Ediacaran Fossils: Bringing Together Palaeontology and Organic Geochemistry | Find, read and cite all the research you need on ResearchGate. Palaeontology and organic geochemistry. Ilya Bobrovskiy, Janet M. Hope, Jochen J. Brocks. Australian National University, Australia. Biological marker molecules; Geochemical fossils; Molecular fossils Biomarkers are the molecular fossils of lipids and other natural products. In sedimentary environments, lipids that escape the... Hydrocarbon biomarkers from Ordovician sediments and the fossil alga *Gloeocapsomorpha prisca* Zalesky 1917. *Geochimica et Cosmochimica Acta*, 51, 2681–2697. CrossRefGoogle Scholar. Holba, A. G., Tegelaar, E. W., Huizinga, B. J., Moldowan, J. M., Singletary, M. S., McCaffrey, M. A., and Dzou, L. I. P., 1998a. Analysis of modern and fossil plant cuticles by curie-point Py-GC and curiepoint Py-GC-MS: recognition of a new highly aliphatic and resistant biopolymer. *Organic Geochemistry*, 10, 769–778. CrossRefGoogle Scholar.